arXiv:math/0412531v2 [math.GT] 30 Dec 2004

Stellar Braiding

Margaret Doig
University of Notre Dame
mdoig@nd.edu

February 1, 2008

Abstract

In this paper, we exhibit an explicit one-dimensional deformation retract $D_k(S_n)$ of the unordered k-point configuration space $U_k^{top}(S_n)$ for any star S_n. These spaces have recently been studied by Abrams and Ghrist. We use this retract to compute an explicit set of free generators β_k of the corresponding braid group $B_k(S_n, c_k)$. In particular, we show that the natural map $i_{k*} : B_{k-1}(S_n, c_{k-1}) \hookrightarrow B_k(S_n, c_k)$ sends β_{k-1} to β_k injectively.

1 Introduction

We investigate the braid group of a star, or the fundamental group of its configuration space. The *unordered k-point configuration space* of a space X is the set of k-element subsets, or

$$U_k^{top}(X) = \{c \subset X : |c| = k\}.$$

This space is topologized as the quotient space of the map

$$p : X^k - \Delta(X) \rightarrow U_k^{top}(X)$$

such that $p(x_1, x_2, \cdots, x_k) = \{x_1, x_2, \cdots, x_k\}$ where the *fat diagonal* of X is

$$\Delta(X) = \{(x_1, x_2, \cdots, x_k) \in X^k : \exists i \neq j \textit{ such that } x_i = x_j\}.$$

The product topology on $X^k - \Delta(X)$ induces a topology on $U_k^{top}(S_n)$.

The *k-point braid group* of a star, $B_k(S_n, c_k)$, is the fundamental group of the configuration space $\pi_1(U_k^{top}(S_n), c_k)$ for some arbitrary basepoint c_k. A *star* S_n is a graph on $n+1$ vertices with one vertex v_0 of degree n and n vertices $v_1, v_2, \cdots, v_n$ of degree one. On S_n, we use the metric

$$d(x, y) = \kappa\rho(x, y) \; \forall x, y \in S_n$$



Figure 1: Vertices in $D_k(S_n)$ are (a) type I if $d(c,v_0)=0$ and (b) type II if $d(c,v_0)=1/2$ (*N.B.:* in (b), $A_1(c)=A_5(c)=A_6(c)=\emptyset$ and so are not arms).

where ρ is the standard simplicial metric on S_n and κ is a fixed constant greater than or equal to $k-1$. Therefore $d(v_0,v_i)=\kappa$ if $i>0$ and $d(v_i,v_j)=2\kappa$ if $1\le i<j\le n$. This induces a Hausdorff metric on the collection of non-empty closed subsets of S_n:

$$d(A,B)=\sup_{\substack{a\in A\\ b\in B}}(d(a,B),d(A,b)).$$

This problem arises in an overlap of topology and graph theory. Graphs treated as topological spaces are well understood, as are configuration spaces of Euclidean spaces. Recently, Abrams and Ghrist introduced the notion of the k-point braid group of a graph (see [3] for an overview of the area). This group is much less clearly understood but is in many ways analogous to Artin's classical braid group, the braid group of the real plane. In the case of a star S, Ghrist has shown that the k-point braid group $B_k(S,c_0)$ is a free group. He has also shown that the associated k-point configuration space $U_k^{top}(\Gamma)$ for any graph Γ is an aspherical space (see [4] Theorem 3.1). Further, he has shown that the natural map $B_{k-1}(\Gamma,c_0)\hookrightarrow B_k(\Gamma,c_0')$ is injective if Γ is a tree (Lemma 2.2).

In Section 2, we exhibit a one-dimensional polyhedron $D_k(S_n)$. We show in Section 3 that the polyhedron is a deformation retract of $U_k^{top}(S_n)$ and name it the *k-spine* of S_n. We then use $D_k(S_n)$ to generate $B_k(S_n,c_k)$ directly. Finally, we show that the natural simplicial map $i_k: D_{k-1}(S_n)\hookrightarrow D_k(S_n)$ induces a monomorphism $i_{k*}: B_{k-1}(S_n,c_{k-1})\hookrightarrow B_k(S_n,c_k)$ mapping a basis β_{k-1} of $B_{k-1}(S_n,c_{k-1})$ into a basis β_k.

2 The space $D_k(S_n)$

We construct a one-dimensional polyhedron $D_k(S_n)$ within $U_k^{top}(S_n)$. Later we will show that it is a deformation retract of $U_k^{top}(S_n)$. Number the vertices $\{v_0,v_1,\cdots,v_n\}$ of S_n

so that $deg(v_0) = n$. The geodesic segment in S_n joining any two points $x,\ y$ is written $[x, y]$. We say a finite subset c of S_n is a *chain* if, for each $x, y \in c$, we have:

$$|c \cap [x, y]| = 1 + d(x, y).$$

Roughly speaking, each point of c is of distance one from its neighbors. We also define:

$$A_i(c) = c \cap [v_0, v_i]$$

for $c \subset S_n$ and call all nonempty $A_i(c)$ the *arms* of c. Then we let a configuration c be *regular* if there is an arm $A_m(c)$ called a *governing arm* where $A_m(c) \cup A_i(c)$ is a chain for all i and there are at least two arms of c. Set

$$D_k(S_n) = \{c : c = regular\}.$$

That is, for any $c \in D_k(S_n)$, there is a governing arm $A_m(c)$ whose union with every other arm is a chain.

If the governing arm is unique, then $0 < d(c, v_0) = d(A_m(c), v_0) < 1/2$; moreover, each arm is a chain, and $d(A_i(c), v_0) = 1 - d(c, v_0)$ for every other arm $A_i(c)$. On the other hand, $A_m(c)$ is not unique exactly when $d(A_i(c), v_0) = 0$ or $1/2$ for each arm $A_i(c)$. In either case, each $c \in D_k(S_n)$ is uniquely determined by the numbers $|A_i(c)|$, by an integer m where $A_m(c)$ is a governing arm, and by the value $d(c, v_0)$. We can now prove the following about $D_k(S_n)$:

Lemma 1. *The space $D_k(S_n)$ is a one-dimensional polyhedron, i.e., a graph.*

Proof. We define a set of vertices of $D_k(S_n)$ by:

$$V(D_k(S_n)) = \{c \in D_k(S_n) : d(c, v_0) = 0 \textit{ or } 1/2\}.$$

There are two types of vertices in $V(D_k(S_n))$. A type I vertex is an element $c \in V(D_k(S_n))$ such that $v_0 \in c$, as in Figure 1a. There is a bicorrespondence between vertices of this type and distributions of $k - 1$ identical particles into the n disjoint edges $(v_0, v_i]$, and so there are $\binom{n+k-2}{n-1}$ of these type I vertices. On the other hand, a vertex c is of type II if $d(c, v_0) = 1/2$ like Figure 2b. There is one of these for each way of distributing k particles in the n edges $(v_0, v_i]$ without putting all the particles on the same edge. Thus, there are $\binom{n+k-1}{n-1} - n$ vertices of this type.

We next define the set of edges of $D_k(S_n)$. Take two vertices c_0 and c_1 such that $d(c_0, c_1) = 1/2$. They will determine an edge of $D_k(S_n)$ denoted $[c_0, c_1]$. Note that exactly one of c_0 and c_1 has type I, say c_0. Then c_1 has type I. There is a unique m such that $|A_m(c_0)| = |A_m(c_1)|$. It follows that $|A_i(c_0)| = |A_i(c_1)| + 1$ for $i \neq m$. Additionally, $d(A_i(c_0), A_i(c_1)) = 1/2$ for each arm $A_i(c_0)$. We may now define the edge $[c_0, c_1]$ of $D_k(S_n)$ by:

$$[c_0, c_1] = \{c \in D_k(S_n) : d(c, c_0) + d(c, c_1) = 1/2\}.$$

Note that a configuration $c \in D_k(S_n)$ is in $[c_0, c_1]$ exactly when $d(A_i(c), A_i(c_0)) \leq 1/2$ and $d(A_i(c), A_i(c_1)) \leq 1/2$. Note that $A_m(c)$ is the unique governing arm of c when $c \neq c_0$ or c_1 and that $|A_i(c)| = |A_i(c_1)|$ for $i \neq m$. The set of edges of $D_k(S_n)$ is:

$$E(D_k(S_n)) = \{[c_0, c_1] : c_0, c_1 \in V(D_k(S_n)) \textit{ and } d(c_0, c_1) = 1/2\}.$$

To complete the proof, we claim that each configuration in $D_k(S_n)/V(D_k(S_n))$ lies in exactly one edge $[c_0, c_1]$ and that each edge $[c_0, c_1]$ is homeomorphic to $[0, 1]$. To see this, let $c \in D_k(S_n)/V(D_k(S_n))$. Recall each configuration is uniquely determined by the numbers $|A_i(c)|$, by an integer m where $A_m(c)$ is a governing arm, and by the value $d(c, v_0)$. Then let c_0 be the type I vertex uniquely determined by $|A_m(c_0)|{=}|A_m(c)|$, by $|A_i(c_0)| - 1 = |A_i(c)|$ for all $i \neq m$. let c_1 be the type II vertex uniquely determined by $|A_i(c_1)| = |A_i(c)|$ for all i. Set $p = d(c, v_0)$. Then $d(c, c_0) = p \leq 1/2$ and $d(c, c_1) = 1/2 - p$. Therefore $c \in [c_0, c_1]$. Additionally, c is uniquely determined by c_0,c_1, and p. A homeomorphism

$$h : [c_0, c_1] \to [0, 1]$$

is then given by the rule $h(c) = 2d(c, v_0)$. □

3 A deformation retraction

The construction of $D_k(S_n)$ suggests a deformation retraction:

Theorem 2. *$D_k(S_n)$ is a deformation retract of $U_k^{top}(S_n)$.*

Proof. Let $c \in U_k^{top}(S_n)$. Without loss of generality, assume $d(A_1(c), v_0) \leq d(A_2(c), v_0) \leq d(A_i(c), v_0)$ for all $i \geq 3$. Set $p = d(A_1(c), v_0)$ and $q = d(A_2(c), v_0)$. We are going to define a homotopy $R : U_k^{top}(S_n) \times [0, 1] \to U_k^{top}(S_n)$ such that $R(c, 0) = c$ and $R(c, 1) = c' \in D_k(S_n)$. If $|A_i(c)| = 0$ for all i, a regular configuration c' is uniquely defined by $|A_i(c')| = |A_i(c)|$ and by $d(A_i(c'), v_0) = 0$ for all i. On the other hand, if $|A_i(c)| \neq 0$ for some i, then $|A_i(c)| \neq 0$ for all i, and then a regular configuration c' is uniquely defined by $|A_i(c')| = |A_i(c)|$ and by $d(A_1(c'), v_0) = \frac{p}{p+q}$ and $d(A_i(c'), v_0) = \frac{q}{p+q}$ for all $i > 1$. There is a natural ordering on the sets $A_i(c)$ and $A_i(c')$ given by $d(x_{ij}, v_0)$ for $x_{ij} \in A_i(c)$ or $A_i(c')$. Then $A_i(c)$ and $A_i(c')$ correspond to points in $[0, k-1] \times [0, k-1] \times \cdots \times [0, k-1]$ below the fat diagonal, and so the straight line segment between them does not cross the fat diagonal and gives a homotopy h_i. These coalesce to provide a homotopy $R : U_k^{top}(S_n) \times [0, 1] \to U_k^{top}(S_n)$. See Figure 2. □

We may now calculate $B_k(S_n, c_0)$ via $\pi_1(D_k(S_n), c_0)$. Let c_0 be the unique vertex contained in $[v_0, v_1]$. Ghrist gives the order of $\pi_1(C_k(S_n), c_0)$, the fundamental group of the ordered configuration space, i.e., of $U_k^{top}(S_n)$ where each element is an ordered n-tuple rather than a set [4]. He uses an inductive proof; we demonstrate the order follows as a corollary of Theorem 2.

Corollary 3. *The braid group $B_k(S_n, c_0)$ is free on $\varphi(k, n)$ generators where*

$$\varphi(k, n) = 1 + (n-1)\binom{n+k-1}{n-1} - \binom{n+k-1}{n-1}.$$

Proof. Note that $D_k(S_n)$ is one-dimensional and so has a free group as fundamental group. The rank can be computed from the Euler characteristic χ of $D_k(S_n)$. We can



Figure 2: Deformation retract R taking $U_5^{top}(S_6)$ to $D_5(S_6)$, mapping (a) configuration c to a type I vertex c' with $d(c', v_0) = 0$ and (b) configuration c to a non-vertex point c' with governing arm $A_4(c')$, that is, with $d(A_4(c'), v_0) = p < 1/2$ and $d(A_i(c'), v_0) = 1-p$ for all i where $A_i(c) \neq \emptyset$ and $A_i(c')$ is an arm.

count the edges in $E(D_k(S_n))$. Since each edge contains exactly one vertex of type I, we need merely add the degrees of the type I vertices. Each of the n vertices c_0 which are contained entirely in one arm $A_m(c_0)$ have degree $n-1$ because each is adjacent to the $n-1$ vertices c_1 where $|A_m(c_1)| = k-1$ and $|A_i(c_1)| = 1$ for $i \neq m$. On the other hand, the remaining $\binom{n+k-1}{n-1} - n$ type I vertices c_0 have degree n because each is adjacent to n different c_1 where $|A_m(c_1)| = |A_m(c_0)|$ for some m and $|A_i(c_1)| = |A_i(c_0)| - 1$ for all $i \neq m$. Thus

$$|E(D_k(S_n))| = n\binom{n+k-1}{n-1} - n.$$

We can also count the vertices since

$$|V(D_k(S_n))| = \binom{n+k-2}{n-1} + \binom{n+k-1}{n-1} - n.$$

Then

$$rank(B_k(S_n)) = 1 - \chi = 1 + (n-1)\binom{n+k-2}{n-1} - \binom{n+k-1}{n-1}.$$

□

Example. *By Corollary 3,*

$$B_k(S_3) \cong \mathcal{F}_{\binom{k}{2}},\ B_2(S_n) \cong \mathcal{F}_{\binom{n-1}{2}},\ \text{and } B_3(S_n) \cong \mathcal{F}_{\frac{2n+3}{3}\binom{n-1}{2}}$$

where $\mathcal{F}_p$ is the free group on p letters. See Figure 3 for the illustrations of $D_k(S_n)$ as generated by the method in the proof of Lemma 1 in these four cases.

Finally, Theorem 2 suggests a monomorphism which maps a basis of $B_{k-1}(S_n, c_{k-1})$ into a basis of $B_k(S_n, c_k)$:

Corollary 4. *Let β_j be a free basis for $B_j(S_n, c_j)$. Then the standard simplicial map $i_k : D_{k-1}(S_n) \hookrightarrow D_K(S_n)$ induces a monomorphism $i_{k*} : B_{k-1}(S_n, c_{k-1}) \hookrightarrow B_k(S_n, c_k)$ which maps β_{k-1} injectively into β_k.*

Proof. Whenever there is a simplicial inclusion $i : L \hookrightarrow M$ for L and M one-dimensional complexes, their fundamental groups $\pi_1(L)$ and $\pi_1(M)$ are free with bases $\beta(L)$ and $\beta(M)$ such that $i_* : \pi_1(L) \hookrightarrow \pi_1(M)$ is injective and $i_*(\beta(L)) \subseteq \beta(M)$. Then $D_{k-1}(S_n)$ and $D_k(S_n)$ are one-dimensional simplicial complexes with a natural embedding $i_k : D_{k-1}(S_n) \hookrightarrow D_k(S_n)$ which maps $c \in D_{k-1}(S_n)$ to $c \cup \{x\}$ for x the unique point such that $c \cup \{x\} \in D_k(S_n)$. (*N.B.:* For a fixed k, such a point always exists by our definition of the metric on S_n and of $D_j(S_n)$.) See Figure 4 for four homeomorphic embeddings of $D_{k-1}(S_n)$ into $D_k(S_n)$. □



Figure 3: Examples of the deformation retracts of (a) $D_2(S_3)$, (b) $D_3(S_3)$, (c) $D_2(S_4)$, and (d) $D_3(S_4)$. The braid groups are free on, respectively, one, three, three, and eleven generators. Solid vertices represent type I vertices and unfilled represent type II. The small diagrams show the corresponding configurations of points in S_n.



Figure 4: Four homeomorphic embeddings of $D_2(S_4)$, denoted by solid edges, into $D_3(S_4)$.

4 Conclusion

In this paper, we demonstrate a one-dimensional deformation retract of the configuration space of a star, $U_k^{top}(S_n)$. This allows us to calculate the braid group $B_k(S_n, c_k)$ of the star and gives a natural inclusion of braid groups $B_{k-1}(S_n, c_{k-1}) \hookrightarrow B_k(S_n, c_k)$ which maps a basis to a basis. In a subsequent paper, the author and Frank Connolly prove that, for any linear tree T, $B_k(T, c_k)$ is a right-angled Artin group.

5 Acknowledgements

This paper was funded by the Provost's Office, Mathematics Department, and Arts and Letters/Science Honors Program, all of the University of Notre Dame. Many thanks are owed to Frank Connolly for his encouragement, support, and constant help.

References

[1] A. Abrams, Configuration spaces and braid groups of graphs. Ph.D thesis, UC Berkeley (2000).

[2] A. Abrams, Configuration spaces of colored graphs. *Dedicated to John Stallings on the ocassion of his 65th birthday. Geom. Dedicata* 92 (2002), 185-194.

[3] A. Abrams, R. Ghrist, Finding topology in a factory: Configuration spaces. *Amer. Math. Monthly* 109 (2002), 140-150.

[4] R. Ghrist, Configuration spaces and braid groups on graphs in robotics. *Knots, braids, and mapping class groups - Papers dedicated to Joan S. Birman (New York, 1998)* 29-40, AMS/IP Stud. Adv. Math., 24, *Amer. Math. Soc., Providence, RI,* 2001.